

A NOTE ON GALOIS MODULES AND THE ALGEBRAIC FUNDAMENTAL GROUP OF PROJECTIVE CURVES

AMÍLCAR PACHECO

ABSTRACT. Let X be a smooth projective connected curve of genus $g \geq 2$ defined over an algebraically closed field k of characteristic $p > 0$. Let G be a finite group, P a Sylow p -subgroup of G and $N_G(P)$ its normalizer in G . We show that if there exists an étale Galois cover $Y \rightarrow X$ with group $N_G(P)$, then G is the Galois group of an étale Galois cover $\mathcal{Y} \rightarrow \mathcal{X}$, where the genus of $\mathcal{X}$ depends on the order of G , the number of Sylow p -subgroups of G and g . Suppose that G is an extension of a group H of order prime to p by a p -group P and X is defined over a finite field $\mathbb{F}_q$ large enough to contain the $|H|$ -th roots of unity. We show that integral idempotent relations in the group ring $\mathbb{C}[H]$ imply similar relations among the corresponding generalized Hasse-Witt invariants.

1. INTRODUCTION

Let X be a smooth projective connected curve of genus $g \geq 2$ defined over an algebraically closed field k of characteristic $p > 0$. Given a finite group G when does there exist an étale Galois cover $Y \rightarrow X$ with group G ?

Grothendieck gave a necessary and sufficient condition for this to take place when $|G|$ is prime to p , [5]. Hasse-Witt theory does the same for a p -group G , [20]. Later, the case of a group G which is the extension of a group H of order prime to p by a p -group P necessary and sufficient conditions for G to be such a Galois group were obtained in [16]. This result was extended for a group G which is the extension of any group H by a p -group P , [3]. The result in [16] relied on the naive representation theoretic approach of [14] as well as on techniques of embedding problems and cohomological dimension. The result in [3] relied on modular representation theory.

One natural question is: how far can this type of approach give information on the construction étale Galois covers? We start by discussing why it was needed in the two latter cases to take an extension of a group by a p -group. Next we consider any finite group G and take P to be a Sylow p -subgroup of G and let $N_G(P)$ be its normalizer in G . Subject to the hypothesis that there exists an étale Galois cover $Y \rightarrow X$ with group $N_G(P)$, we show that G is the Galois group of an étale Galois cover $\mathcal{Y} \rightarrow \mathcal{X}$ (using patching theory [19]), where the genus of $\mathcal{X}$ depends on the order of G , the number of Sylow p -subgroups of G and g .

Finally, the necessary and sufficient condition of [16] can be expressed in terms of certain numbers, called generalized Hasse-Witt invariants, which generalize the notion of the p -rank of a curve. Our next result is of a different nature, however

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* 14H30, 14G15.

This work was partially supported by CNPq research grant 300896/91-3 and Pronex #41.96.0830.00.

related to these numbers. At this point we assume that H is a group of order prime to p and the X is defined over a finite field $\mathbb{F}_q$ large enough to contain the $|H|$ -th roots of unity. We show that integral idempotent relations in the group ring $\mathbb{C}[H]$ imply similar relations among the corresponding generalized Hasse-Witt invariants. Analogous situations for the genera and Hasse-Witt invariants of curves were done before in [15] and [10]. As in [15], where relations among zeta-functions are first proved, the result follows from relations between Artin L -functions.

2. GALOIS GROUPS OF ÉTALE GALOIS COVERS

Let X be a smooth projective connected curve of genus $g \geq 2$ defined over an algebraically closed field k of characteristic $p > 0$. Let $\pi_1(X)$ be its algebraic fundamental group. Let Γ_g be the topological fundamental group of a compact Riemann surface of genus g and $\hat{\Gamma}_g$ its profinite completion. It is a result due to Grothendieck [5, Corollary 2.12] that there exists a surjective group homomorphism $\varphi : \hat{\Gamma}_g \twoheadrightarrow \pi_1(X)$ whose kernel is contained in every open normal subgroup N of $\hat{\Gamma}_g$ such that $(\hat{\Gamma}_g : N)$ is prime to p . In particular, $\pi_1(X)$ is topologically finitely generated. Let $\pi_A(X)$ be the set of isomorphism classes of finite groups which are quotients of $\pi_1(X)$, i.e., $G \in \pi_A(X)$ if and only if there exists an étale Galois cover $Y \rightarrow X$ such that $G = \text{Gal}(Y/X)$. Hence, the determination of $\pi_1(X)$ is equivalent to that of $\pi_A(X)$.

Our first goal is to discuss $\pi_A(X)$. As a consequence of [5, Corollary 2.12] Grothendieck obtained that a finite group G of order prime to p lies in $\pi_A(X)$ if and only if G is isomorphic to a finite quotient of $\hat{\Gamma}_g$. The case of finite p -groups P is first treated by Hasse-Witt theory, when P is a p -elementary abelian group [20, §11]. In this case, $G \in \pi_A(X)$ if and only if it has at most γ_X generators, where γ_X is the $\mathbb{F}_p$ -dimension of the p -torsion subgroup $J_X[p]$ of the Jacobian variety J_X of X . This number is called the Hasse-Witt invariant or the p -rank of X . The general case is reduced to this one by taking the quotient $P/\Phi(P)$ of P by its Frattini subgroup $\Phi(P) = [P, P]P^p$ and using that $\pi_1(X)$ has p -cohomological dimension $\text{cd}_p(\pi_1(X))$ at most 1 [16, Introduction].

The first mixed case to consider is that of a finite group G which is an extension of a group H of order prime to p by a p -group P [16, Theorem 1.3]. We proved that the necessary and sufficient condition for $G \in \pi_A(X)$ is that there exists an étale Galois cover $Y \rightarrow X$ such that $P/\Phi(P)$ injects in $J_Y[p]$ as an $\mathbb{F}_p[H]$ -module. Moreover, the étale Galois cover $Z \rightarrow X$ with group G dominates $Y \rightarrow X$. In the next section we will express this explicitly in terms of multiplicities of irreducible representations. It should be also noted that in [19, Corollary 3.5], using formal patching, it was proved that a finite group G having at most g generators lies in $\pi_A(\mathcal{X})$, for some smooth projective connected curve $\mathcal{X}$ of genus g . In contrast, the first step of the proof of the former result is based in the case where P is a p -elementary abelian group and it was solved by naive representation theory in [14, Propositions 2.4 and 2.5]. We will comment later on how to go from this step to the general case.

Later Borne in [3, Theorem 1.1] used modular representation theory to extend [16, Theorem 1.3] to the case of a finite group G which is the extension of any group H by a p -group P . He also obtained a necessary and sufficient condition for $G \in \pi_A(X)$ in terms of representation theory. At this point it is convenient to phrase these two latter results in terms of embedding problems [6, p. 366]. Let $\Pi, \mathcal{G}, \mathcal{H}$ and

$\mathcal{P}$ be profinite groups. An embedding problem is a pair $\mathcal{E} = (\mathcal{G} \twoheadrightarrow \mathcal{H}, \Pi \twoheadrightarrow \mathcal{H})$ of surjective profinite group homomorphisms. We say that $\mathcal{E}$ has a weak, respectively strong, solution if there exists a group homomorphism, respectively a surjective group homomorphism, $\Pi \rightarrow \mathcal{G}$ such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{G} & \twoheadrightarrow & \mathcal{H} \\ & \nwarrow & \uparrow \\ & & \Pi \end{array}$$

The two latter situations can be described as obtaining a strong solution to the embedding problem

$$\begin{array}{ccccccc} 0 & \longrightarrow & P & \longrightarrow & G & \longrightarrow & H \longrightarrow 0 \\ & & & & & \nwarrow & \uparrow \\ & & & & & & \pi_1(X) \end{array}$$

The kernel $\mathcal{P}$ of $\mathcal{G} \twoheadrightarrow \mathcal{H}$ is called the kernel of the embedding problem $\mathcal{E}$. In the case where $\mathcal{P}$ is a profinite p -group, we say that $\mathcal{E}$ is a p -embedding problem [7, Section 2]. Thus, [16, Theorem 1.3] and [3, Theorem 1.1] give a necessary and sufficient condition for finite p -embedding problems for the fundamental group of projective curves to be strongly solved.

One natural question is: how far can the Galois module theory approach go in order to obtain information on $\pi_A(X)$? Let us first point out where the p -group kernel was used in both situations. In [16, §6], we started from the strong solution of the embedding problem $(\pi_1(X) \twoheadrightarrow H, P/\Phi(P) \rtimes H \twoheadrightarrow H)$. This solution was obtained via naive representation theory, [14, Propositions 2.4 and 2.5]. Then we used the p -group kernel and $\text{cd}_p(\pi_1(X)) \leq 1$ to obtain via Galois cohomology [21, Proposition 16, I-23] a weak solution to the embedding problem $(\pi_1(X) \twoheadrightarrow P/\Phi(P) \rtimes H, G \twoheadrightarrow P/\Phi(P) \rtimes H)$. The strong solution is obtained via a Frattini type of argument, [16, Remark 4.4]. In [3, §2] the p -group kernel was used to obtain (via Clifford's theorem) a 1-1 correspondence between the set of isomorphism classes of simple $k[H]$ -modules and the set of isomorphism classes of simple $k[G]$ -modules [3, Remark 2 to Proposition 2.4]. This was crucial in the proof of the reduction lemma [3, Lemma 3.1], which was used to prove [3, Theorem 1.1].

Abhyankar's conjecture is closely related to the problem of when a finite group G lies in $\pi_A(X)$, [1]. We now describe this conjecture. Let $x_1, \dots, x_r \in X$ with $r \geq 1$ and $U = X - \{x_1, \dots, x_r\}$. Let $\pi_1(U)$ be the algebraic fundamental group of U and $\pi_A(U)$ the set of isomorphism classes of finite groups which are quotients of $\pi_1(U)$. The conjecture states that a finite group $G \in \pi_A(U)$ if and only if $G/p(G) \in \pi_A(U)$, where $p(G)$ is the quasi- p subgroup of G , i.e., the group generated by the Sylow p -subgroups of G . One side of this conjecture was already known to Grothendieck. In fact, another consequence of [5, Corollary 2.12] is that a finite group G of order prime to p lies in $\pi_A(U)$ if and only if G is a finite quotient of the profinite completion $\hat{\Gamma}_{g,r}$ of the topological fundamental group $\Gamma_{g,r}$ of a compact Riemann surface of genus g minus r points. If $G \in \pi_A(U)$, then $G/p(G) \in \pi_A(U)$ and Grothendieck's result give a necessary and sufficient condition for this to take place. The converse was the issue in the conjecture. When U is the affine line, the conjecture was proved by Raynaud [17], using rigid analytic techniques. Later, Harbater [8] extended this result to any affine curve using formal geometry.

The analogue of Abhyankar's conjecture fails for projective curves. Indeed, suppose X is an ordinary curve, i.e., $J_X[p] \cong (\mathbb{Z}/p\mathbb{Z})^g$. Let $n > g$ be an integer and $G = (\mathbb{Z}/p\mathbb{Z})^n$. Then $G/p(G) = \{1\} \in \pi_A(X)$, however $G \notin \pi_A(X)$, since it violates the condition that the number of generators of G has to be at most $\gamma_X = g$. The representation theoretic conditions of [16, Theorem 1.3] and [3, Theorem 1.1] are the obstruction for the condition $G/p(G) \in \pi_A(X)$ to be sufficient for $G \in \pi_A(X)$.

The first idea to proceed is to first obtain a strong solution to the embedding problem $(\pi_1(X) \twoheadrightarrow H, (p(G)/[p(G)]) \rtimes H \twoheadrightarrow H)$ in a similar way to [14, Propositions 2.4 and 2.5], where $[p(G)]$ denotes the commutator of $p(G)$. Then we are lead to try to solve the embedding problem $(\pi_1(X) \twoheadrightarrow (p(G)/[p(G)]) \rtimes H, G \twoheadrightarrow (p(G)/[p(G)]) \rtimes H)$. However, the kernel is not any longer a p -group and the ℓ -cohomological dimension of $\pi_1(X)$ is not necessarily 1, for a prime number $\ell \neq p$ (cf. [2, Corollaire 4.3] and [16, §5]). In the modular representation theory approach, although [3, Proposition 2.4] can be proved without the hypothesis of the embedding problem having a p -kernel, [3, Lemma 3.1] cannot. So this points to a bound on the Galois module theoretic method.

The strategy of strongly solving first a finite p -embedding problem for the affine line $\mathbb{A}^1$ in order to show that $G/p(G) \in \pi_A(\mathbb{A}^1)$ implies $G \in \pi_A(\mathbb{A}^1)$ appears in [17]. There rigid patching is used together with other techniques. Strongly solving finite p -embedding problems also appears as part of the strategy of [8], where also a prescribed local behavior is needed [7] and formal patching replaces rigid patching. For an account of the sketch of both proofs see [9, p. 85, 86].

If we try to mimic the use of patching theory for projective curves, not only do we not realize G as a Galois group of an étale Galois cover of X , but also the base curve has its genus increased. This is due to [19, Theorem 3.1] which we describe next.

Given an integer $g \geq 1$, let $\pi_A(g)$ be the set of isomorphism classes of finite groups G such that $G \in \pi_A(X)$, for some smooth projective connected curve X of genus g defined over k . Our main ingredient is the following result.

Proposition 2.1. [19, Theorem 3.1] *Let G be a finite group generated by two subgroups H_1 and H_2 . Suppose there exist smooth projective connected curves X_1 and X_2 of genus g_1 and g_2 , respectively, defined over k such that $H_1 \in \pi_A(U_1)$ and $H_2 \in \pi_A(U_2)$, where $U_1 = X_1 - \{\xi_1\}$ and $U_2 = X_2 - \{\xi_2\}$. Suppose furthermore that the ramification over ξ_1 and ξ_2 is tame and that the generator of the inertia group of ξ_1 is the inverse in G of that of the inertia group at ξ_2 . Then $G \in \pi_A(g_1 + g_2)$.*

Let P be a Sylow p -subgroup of G and $N_G(P)$ its normalizer in G . Let $|G| = mp^e$ with $p \nmid m$ and n_p the number of Sylow p -subgroups of G . By [4, Chapter I, Theorem 3.7] $G = N_G(P)p(G)$. In particular, G is generated by $N_G(P)$ and $p(G)$.

Theorem 2.2. *Suppose there exists a smooth projective connected curve X of genus $g \geq 2$ such that $N_G(P) \in \pi_A(X)$ (hence satisfying the condition of [16, Theorem 1.3]). Then $G \in \pi_A(n_p + m(g-1) + g)$.*

Proof. Let $Z(P) \rightarrow X$ be an étale Galois cover with group $N_G(P)$. This cover dominates an étale Galois cover $Y(P) \rightarrow X$ with group $N_G(P)/P$ and $P \in \pi_A(Y(P))$. By the Riemann-Hurwitz formula $Y(P)$ has genus $(m/n_p)(g-1) + 1$. Note that for any other Sylow p -subgroup P' of G , $N_G(P') \cong N_G(P)$, hence $N_G(P') \in \pi_A(X)$. Consequently, there exists an étale Galois cover $Z(P') \rightarrow X$ with group $N_G(P')$ which dominates an étale Galois cover $Y(P') \rightarrow X$ with group $N_G(P')/P'$ and

$P' \in \pi_A(Y(P'))$. By Proposition 2.1, since $p(G)$ is generated by the Sylow p -subgroups of G , $p(G) \in \pi_A(m(g-1) + n_p)$. Let $\mathcal{Y}' \rightarrow \mathcal{X}'$ be an étale Galois cover with group $p(G)$. Since $N_G(P) \in \pi_A(X)$, i.e., $N_G(P)$ is the Galois group of an étale Galois cover $Y \rightarrow X$, once again applying Proposition 2.1, we conclude $G \in \pi_A(n_p + m(g-1) + g)$. $\square$

3. IDEMPOTENT RELATIONS AND GENERALIZED p -RANKS OF CURVES

Let G be a finite group which is the extension of a group H of order prime to p by a p -group P . Let $\Phi(P) = [P, P]P^p$ be the Frattini subgroup of P . Let X be a smooth projective connected curve of genus $g \geq 2$ over k . The necessary and sufficient condition for G to lie in $\pi_A(X)$ can be phrased in an explicit way using representation theory.

For each irreducible character χ of H , let V_χ be the corresponding irreducible $k[H]$ -module and $\mathbb{F}_{p^{\iota_\chi}} = \mathbb{F}_p(\chi)$ the field generated by $\mathbb{F}_p$ and the values of χ . Let ρ_χ be the irreducible representation corresponding to V_χ . Then for each positive power p^m of p we have another irreducible representation $\rho_{\chi^{p^m}}$ of G such that $\rho_{\chi^{p^m}}(\tau) = \rho_\chi(\tau)^{p^m}$, for every $\tau \in H$. Denote by $V_{\chi^{p^m}}$ the irreducible $k[G]$ -module corresponding to $\rho_{\chi^{p^m}}$. The irreducible $\mathbb{F}_p[H]$ -modules are given by $V_{[\chi]} = V_\chi \oplus V_{\chi^p} \oplus \dots \oplus V_{\chi^{p^{\iota_\chi}-1}}$. Let $P/\Phi(P) = \bigoplus_{[\chi]} V_{[\chi]}^{m_{[\chi]}}$ be the decomposition of $P/\Phi(P)$ into irreducible $\mathbb{F}_p[H]$ -modules.

Let $Y \rightarrow X$ a Galois cover with group H . The group H acts on $J_Y[p]$. Let $J_Y[p] \otimes_{\mathbb{F}_p} k = \bigoplus_\chi V_\chi^{r_\chi}$ be the decomposition of $J_Y[p] \otimes_{\mathbb{F}_p} k$ into irreducible $k[H]$ -modules and denote $J_Y[p]_\chi = V_\chi^{r_\chi}$. This is a k -vector space whose dimension $\gamma_{Y,\chi}$ is called the generalized Hasse-Witt invariant of Y with respect to χ [18, §2]. It follows from [14, Proposition 2.8 and Lemma 2.11] that $\gamma_{Y,\chi^{p^m}} = \gamma_{Y,\chi}$ for every positive power p^m of p . Thus, we denote it by $\gamma_{Y,[\chi]}$. The condition, $P/\Phi(P)$ injects as an $\mathbb{F}_p[H]$ -module in $J_Y[p]$ is equivalent to $m_{[\chi]}n_{[\chi]} \leq \gamma_{Y,[\chi]}$ for any irreducible $\mathbb{F}_p[H]$ -module $V_{[\chi]}$, where $n_{[\chi]}$ denotes the dimension of any of the $k[H]$ -irreducible modules $V_{\chi^{p^m}}$. Hence, our representation theoretic condition is expressed in terms of the generalized Hasse-Witt invariants.

These invariants are also used to count the number of étale Galois covers of X whose Galois group G is the extension of a group H of order prime to p by an elementary p -abelian group P such that the action of H on P is faithful and irreducible [14, Theorem 1.8] (for previous results see also [13] and [11]). When instead of an algebraically closed field k we consider a finite $\mathbb{F}_q$ large enough to contain the $|H|$ -th roots of unity, the generalized Hasse-Witt invariants are obtained as the degrees of the L -functions $L(t, \chi, Y/X)^{n_\chi}$ which we now describe [18, Corollary 4.1].

We suppose from now on that Y and X are defined over a finite field $\mathbb{F}_q$ of characteristic p large enough to contain the $|H|$ -th roots of unity. Let $F : Y \rightarrow Y$ be the geometric Frobenius morphism of Y with respect to $\mathbb{F}_q$. For each positive integer ν and for each $\tau \in H$, let $\Lambda(\tau \circ F^\nu)$ be the number of points of Y fixed by $\tau \circ F^\nu$. The Artin L -function $L(t, \chi, Y/X)$ of an irreducible character χ of H is defined as

$$L(t, \chi, Y/X) = \exp \left(\sum_{\nu \geq 1} A_\nu(\chi) \frac{t^\nu}{\nu} \right), \text{ where } A_\nu(\chi) = \frac{1}{|H|} \sum_{\tau \in H} \chi(\tau^{-1}) \Lambda(\tau \circ F^\nu).$$

Let $\ell \neq p$ be a prime number and let $T_\ell(J_Y) = \varprojlim J_Y[\ell^n]$ be the ℓ -adic Tate module of J_Y . It is a result due to Weil [12, p. 186, Theorem 6] that there exists a representation $\text{End}(J_Y) \rightarrow \text{End}(T_\ell(J_Y))$ given by $\tau \mapsto \alpha_\tau^*$. Let ϑ be the restriction of this representation to H . Denote also by F the geometric Frobenius of J_Y over $\mathbb{F}_q$ and let $\alpha_{F^\nu}^* \in \text{End}(T_\ell(J_Y))$ be the element corresponding to ν -th power F^ν of F . Then $\Lambda(\tau \circ F^\nu) = q^\nu + 1 - \text{Tr}(\alpha_{\tau^{-1}}^* \circ \alpha_{F^\nu}^* | T_\ell(J_Y))$. Hence,

$$L(t, \chi, Y/X) = \frac{P(t, \chi, Y/X)}{(1 - qt)^{u_\chi} (1 - t)^{u_\chi}},$$

where

$$(3.1) \quad P(t, \chi, Y/X) = \exp \left(\sum_{\nu \geq 1} -\frac{1}{|H|} \left(\sum_{\tau \in H} \chi(\tau^{-1}) \text{Tr}(\alpha_{\tau^{-1}}^* \circ \alpha_{F^\nu}^*) \right) \frac{t^\nu}{\nu} \right)$$

is a polynomial in t and $u_\chi = 1$, if χ is trivial, $u_\chi = 0$, otherwise. Denote by χ^{-1} the character of the representation $\rho_\chi(\tau)^{-1}$, for every $\tau \in H$.

For each subgroup M of H and each irreducible character χ_M of M , let $e_{\chi_M} = \frac{\chi_M(1)}{|M|} \sum_{\sigma \in M} \chi_M(\sigma^{-1}) \sigma \in \mathbb{C}[H]$ be the corresponding idempotent. An integral idempotent relation is an equality of the form $\sum_M s_{\chi_M} e_{\chi_M} = 0$, where M runs through the subgroups of H , $s_{\chi_M} \in \mathbb{Z}$ and for each M we choose an irreducible character χ_M of M .

Theorem 3.1. *An integral idempotent relation $\sum_M s_{\chi_M} e_{\chi_M} = 0$ in $\mathbb{C}[H]$ implies a relation*

$$\prod_M P(t, \chi_M^{-1}, Y/Y_M)^{\chi_M(1) s_{\chi_M}} = 1,$$

where Y_M is the quotient curve Y/M and $Y \rightarrow Y_M$ the corresponding Galois cover.

Proof. Let $\mathbb{C}_\ell$ be the completion of the algebraic closure $\overline{\mathbb{Q}_\ell}$ of $\mathbb{Q}_\ell$ and identify the $|H|$ -th roots of unity in $\mathbb{C}$ with those in $\mathbb{C}_\ell$. We will consider the integral idempotent relation $\sum_M s_{\chi_M} e_{\chi_M} = 0$ inside $\mathbb{C}_\ell[H]$ instead of $\mathbb{C}[H]$. Extend the representation ϑ to a representation $\overline{\vartheta} : \mathbb{C}_\ell[H] \rightarrow \text{GL}(T_\ell(J_Y) \otimes_{\mathbb{Z}_\ell} \mathbb{C}_\ell)$. Then

$$0 = \overline{\vartheta} \left(\sum_M s_{\chi_M} e_{\chi_M} \right) = \sum_M \left(\frac{s_{\chi_M} \chi_M(1)}{|M|} \right) \sum_{\sigma \in M} \chi_M(\sigma^{-1}) \alpha_\sigma^*.$$

Taking the trace of following representation in $T_\ell(J_Y) \otimes_{\mathbb{Z}_\ell} \mathbb{C}_\ell$, we obtain

$$0 = \text{Tr} \left(\overline{\vartheta} \left(\sum_M s_{\chi_M} e_{\chi_M} \right) \circ \alpha_{F^\nu}^* \right) = \sum_M \frac{s_{\chi_M}}{|M|} \sum_{\sigma \in M} \chi_M(\sigma^{-1}) \text{Tr}(\alpha_\sigma^* \circ \alpha_{F^\nu}^*).$$

The result now follows from (3.1). $\square$

The Artin L -function $L(t, Y/X, \chi)$ can be expressed p -adically as follows. For each $P \in J_Y$, let $\mathcal{O}_{J_Y, P}$ be the local ring of J_Y at P . For each integer $n \geq 1$, let $W_n(\mathcal{O}_{J_Y, P})$ be the Witt ring of length n with coefficients in $\mathcal{O}_{J_Y, P}$. Let $\mathcal{W}_n$ be the sheaf over J_Y defined by these rings. The cohomology groups $H^m(J_Y, \mathcal{W}_n)$ form a projective limit. Let $H^m(J_Y, \mathcal{W}) = \varprojlim H^m(J_Y, \mathcal{W}_n)$. Let J_Y^Δ be the dual abelian variety of J_Y and $T_p(J_Y^\Delta)$ its p -adic Tate module. Let $W(\overline{\mathbb{F}_q})$ the ring of Witt vectors of $\overline{\mathbb{F}_q}$ and $\overline{T}_p(J_Y^\Delta) = T_p(J_Y^\Delta) \otimes_{\mathbb{Z}_p} W(\overline{\mathbb{F}_q})$. Let $L(J_Y) = H^1(J_Y, \mathcal{W}) \oplus \overline{T}_p(J_Y^\Delta)$. Note that identifying the $|H|$ -th roots of unity in $\mathbb{C}$ with those in $\mathbb{F}_q$, we can assume that the characters of H take value in $\mathbb{F}_q$, hence in $W(\overline{\mathbb{F}_q})$. Let $L(J_Y) = \bigoplus_\chi \mathcal{V}_\chi^{m_\chi}$ be the

decomposition of $L(J_Y)$ into irreducible $W(\overline{\mathbb{F}}_q)[H]$ -modules and let $L(J_Y)_\chi = \mathcal{V}_\chi^{m_\chi}$. It is proved in [18, Corollary 3.1] that $P(t, \chi, Y/X)^{\chi(1)} = \det(1 - \alpha_{p,F}^* | L(J_Y)_{\chi^{-1}})$, where $\alpha_{p,F}^* \in L(J_Y)$ is the endomorphism corresponding to the geometric Frobenius F of J_Y . In particular, $P(t, \chi, Y/X)$ has p -integral coefficients.

Let $K = \mathbb{F}_q(Y)$, let Ω_K be the space of all differentials of K and $H^0(Y, \Omega_Y)$ the subspace of regular differentials. Let $u \in K$ be a separating variable and $\omega = fdu \in \Omega_K$. The Cartier operator is defined by $\mathcal{C}(\omega) = (-d^{p-1}f/du^{p-1})^{1/p}du$. It is a $1/p$ -linear operator, i.e., $\mathcal{C}(a^p\omega) = a\omega$, for $a \in K$, and it acts on $H^0(Y, \Omega_Y)$ [20, §10, p.39].

Let $H^0(Y, \Omega_Y) = \bigoplus_\chi H^0(Y, \Omega_Y)_\chi$ be the decomposition of $H^0(Y, \Omega_Y)$ into a sum of irreducible $\mathbb{F}_q[H]$ -modules, where $H^0(Y, \Omega_Y)_\chi = V_\chi^{m_\chi}$ and V_χ is the irreducible $\mathbb{F}_q[H]$ -module corresponding to χ . Let $q = p^n$, then $\mathcal{C}^n$ acts on $H^0(Y, \Omega_Y)_\chi$. Moreover, $\gamma_{Y,\chi} = \dim_{\mathbb{F}_q} \ker(1 - \mathcal{C}^n | H^0(Y, \Omega_Y)_\chi)$ [18, Proposition 2.3]. It is also shown that $\det(1 - \alpha_{p,F}^* t | L(J_Y)_{\chi^{-1}}) \pmod{p} = \det(1 - \mathcal{C}^n t | H^0(Y, \Omega_Y)_\chi)$ [18, Theorem 4.1]. Consequently, $\deg(\det(1 - \alpha_{p,F}^* t | L(J_Y)_{\chi^{-1}}) \pmod{p}) = \gamma_{Y,\chi}$ and we obtain the following corollary of Theorem 3.1.

Corollary 3.2. *An integral idempotent relation $\sum_M s_{\chi_M} e_{\chi_M} = 0$ in $\mathbb{C}[H]$ implies a relation*

$$\sum_M s_{\chi_M} \gamma_{Y, \chi_M^{-1}} = 0.$$

REFERENCES

- [1] S. Abhyankar, *Coverings of algebraic curves*, Amer. J. Math. **78** (1957), 825-856.
- [2] M. Artin, *Dimension cohomologique: premiers résultats*, SGA 4, Lecture Notes in Math. **269**, 43-63.
- [3] N. Borne, *A relative Shafarevich theorem*, preprint, 2001.
- [4] D. Gorenstein, *Finite Groups*, Chelsear, 1980.
- [5] A. Grothendieck, *Rêtements étales et groupe fondamental* SGA 1, Lecture Notes in Math. **244**, Springer-Verlag, 1971.
- [6] D. Harbater, *Fundamental groups and embedding problems in characteristic p* , Contemp. Math. **186** (1995), 353-369.
- [7] D. Harbater, *Embedding problems with local conditions*, Israel J. Math. **118** (2000), 317-355.
- [8] D. Harbater, *Abhyankar's conjecture on Galois groups over curves*, Invent. Math. **117** (1994), 1-25.
- [9] D. Harbater, *Patching and Galois theory*, preprint.
- [10] E. Kani, *Relations between the genera and Hasse-Witt invariants of Galois coverings of curves*, Canadian Math. Bulletin **28** (1985), 321-327.
- [11] H. Katsurada, *On generalized Hasse-Witt invariants and unramified Galois extensions of an algebraic function field*, J. Japan Math. Soc. **31** (1979), 101-125.
- [12] S. Lang, *Abelian Varieties*, Springer-Verlag, 1983.
- [13] S. Nakajima, *On generalized Hasse-Witt invariants of an algebraic curve*, Adv. Stud. Pure Math. **2** (1983), 69-88.
- [14] A. Pacheco, *Unramified Galois coverings of algebraic curves*, J. Number Theory **53** (1995), 211-228.
- [15] A. Pacheco, *A note on relations between the zeta-functions of Galois coverings of curves over finite fields*, Canadian Math. Bulletin **33** (1990), 282-285.
- [16] A. Pacheco, K. Stevenson, *Finite quotients of the algebraic fundamental group of projective curves in positive characteristic*, Pacific J. Math **192** (2000), 143-158.
- [17] M. Raynaud, *Rêtements de la droite affine en caractéristique $p > 0$ et conjecture d'Abhyankar*, Invent. Math **116** (1994), 425-462.
- [18] H.-G. Rück, *L -series and class groups of function fields*, J. Number Theory **22** (1986), 177-189.

- [19] K. Stevenson, *Galois groups of unramified covers of projective curves in characteristic p* , J. Algebra **182** (1996), 770-804.
- [20] J.-P. Serre, *Sur la topologie des variétés algébriques en caractéristique p* , in “Symp. Topologia Algebraica (Mexico City), 1956”, 24-53.
- [21] J.-P. Serre, *Cohomologie Galoisienne*, Lecture Notes in Math **5**, 1986, Springer-Verlag.

UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, DEPARTAMENTO DE MATEMÁTICA PURA, RUA
GUAIAQUIL 83, CACHAMBI, 20785-050 RIO DE JANEIRO, RJ, BRASIL
E-mail address: `amilcar@impa.br`